

Received: 15 April 2018

Accepted: 27 June 2018




DOI: 10.1111/1365-2745.13039

RESEARCH ARTICLE

Journal of Ecology



Globally consistent impact of tropical cyclones on the structure of tropical and subtropical forests

Thomas Ibanez^{1,2}  | Gunnar Keppel^{3,4}  | Christophe Menkes⁵ | Thomas W. Gillespie⁶ |
Matthieu Lengaigne^{7,8} | Morgan Mangeas⁹ | Gonzalo Rivas-Torres^{10,11}  |
Philippe Birnbaum^{1,2,12}

¹Institut Agronomique néo-Calédonien (IAC), Equipe Sol & Végétation (SolVeg), Nouméa, New Caledonia; ²AMAP, Univ Montpellier, CIRAD, CNRS, INRA, IRD, Montpellier, France; ³School of Natural and Built Environments and Future Industries Institute, University of South Australia, Adelaide, South Australia, Australia; ⁴Biodiversity, Macroecology and Biogeography Group, Faculty of Forest Sciences and Forest Ecology, University of Göttingen, Göttingen, Germany; ⁵IRD/Sorbonne Universités (UPMC, Université Paris 06)/CNRS/MNH, LOCEAN – UMR 7159, Nouméa, New Caledonia; ⁶Department of Geography, University of California Los Angeles, Los Angeles, California; ⁷Sorbonne Universités (UPMC, Univ Paris 06)-CNRS-IRD-MNH, LOCEAN Laboratory, IPSL, Paris, France; ⁸Indo-French Cell for Water Sciences, IISc-NIO-IITM-IRD Joint International Laboratory, NIO, Goa, India; ⁹UMR ESPACE-DEV, IRD, Univ. La Réunion, Univ. Guyane, Univ. Antilles, Univ. Montpellier, Montpellier, France; ¹⁰Colegio de Ciencias Biológicas y Ambientales and Galápagos Academic Institute for the Arts and Sciences, Universidad San Francisco de Quito, Quito, Ecuador; ¹¹Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, Florida and ¹²CIRAD, UMR AMAP, Nouméa, Nouvelle-Calédonie

Correspondence

Thomas Ibanez, Institut Agronomique néo-Calédonien, Equipe SolVeg, BPA5, 98800 Nouméa, New Caledonia.
Email: ibanez@iac.nc

Funding information

Agency for Economic and Environmental Development (DDEE) of the North Province of New Caledonia

Handling Editor: Emily Lines

Abstract

1. Tropical cyclones (TCs) are large-scale disturbances that regularly impact tropical forests. Although long-term impacts of TCs on forest structure have been proposed, a global test of the relationship between forest structure and TC frequency and intensity is lacking. We test on a pantropical scale whether TCs shape the structure of tropical and subtropical forests in the long term.
2. We compiled forest structural features (stem density, basal area, mean canopy height and maximum tree size) for plants ≥ 10 cm in diameter at breast height from published forest inventory data (438 plots ≥ 0.1 ha, pooled into 250 1×1 -degree grid cells) located in dry and humid forests. We computed maps of cyclone frequency and energy released by cyclones per unit area (power dissipation index, PDI) using a high-resolution historical database of TCs trajectories and intensities. We then tested the relationship between PDI and forest structural features using multivariate linear models, controlling for climate (mean annual temperature and water availability) and human disturbance (human foot print).
3. Forests subject to frequent cyclones (at least one TCs per decade) and high PDI exhibited higher stem density and basal area, and lower canopy heights. However, the relationships between PDI and basal area or canopy height were partially masked by lower water availability and higher human foot print in tropical dry forests.
4. *Synthesis.* Our results provide the first evidence that tropical cyclones have a long-term impact on the structure of tropical and subtropical forests in a globally consistent way. The strong relationship between power dissipation index and stem density suggests that frequent and intense tropical cyclones reduce canopy cover

through defoliation and tree mortality, encouraging higher regeneration and turnover of biomass. The projected increase in intensity and poleward extension of tropical cyclones due to anthropogenic climate change may therefore have important and lasting impacts on the structure and dynamics of forests in the future.

KEYWORDS

basal area, canopy height, hurricanes, power dissipation index, stem density, tropical dry forest, tropical humid forest, typhoons

1 | INTRODUCTION

Tropical cyclones (TCs), also referred to as hurricanes in the Atlantic and Northeast Pacific and typhoons in the Northwest Pacific, are unavoidable, large-scale disturbances (Everham & Brokaw, 1996; Turner, Baker, Peterson, & Peet, 1998; Turner & Dale, 1998; Whigham, Dickinson, & Brokaw, 1999). Disturbances are “relatively discrete events in time that disrupt ecosystem, community, or population structure and change resource or substrate availability or the physical environment” (White & Pickett, 1985). Due to their large footprints (TCs signatures extend over hundreds of kilometres and their tracks cover thousands of km, see Chan & Chan, 2015; Knaff, Longmore, & Molenar, 2014) and high intensity (sustained wind speeds of up to 300 km/hr), TCs can cause extensive damages to natural ecosystems and human societies (Costanza et al., 2008;

Lugo, 2008; Yih, Boucher, Vandermeer, & Zamora, 1991). This was recently highlighted by the devastating impacts of Cyclone Winston (2016), Hurricane Irma (2017) and Typhoon Haiyan (2013), some of the most intensive systems ever recorded (Le Page, 2016).

The effects of disturbances on the composition, structure and functioning of ecosystems depend on the disturbance regime, which is for TCs mainly characterized by their frequency, size and intensity (Turner et al., 1998; Whelan, 1995; White & Jentsch, 2001; White & Pickett, 1985). The disturbance regime exerted by TCs is highly spatially heterogeneous. For example, forests in the Northwest Pacific and Northwest Atlantic experience frequent, intense and large TCs while forests in South America and Africa experience few TCs (Chan & Chan, 2015; Schreck, Knapp, & Kossin, 2014). Therefore, on a pantropical scale forests may experience TC disturbance regimes ranging from infrequent TCs to frequent and intense TCs (Figure 1).




FIGURE 1 Global distribution of the frequency and energy released (power dissipation index) by tropical cyclones, and locations of forest structure datasets [Colour figure can be viewed at wileyonlinelibrary.com]

On a landscape scale, the TC impacts vary with topographical exposure and other abiotic and biotic factors (e.g., Basnet, Likens, Scatena, & Lugo, 1992; Boose, Serrano, & Foster, 2004; Everham & Brokaw, 1996; Metcalfe, Bradford, & Ford, 2008; Turton, 2008; Webb, Seamon, & Fa'aumu, 2011). However, in regions where TCs are frequent, successive TCs with different trajectories and wind directions should in the long term affect the entire landscape (Turner, Gardner, & O'Neill, 2001).

TCs can considerably impact the structure of forests (Everham & Brokaw, 1996; Turner, Dale, & Everham, 1997; Whigham et al., 1999). Visible short- to medium-term effects (Lugo, 2008) include direct damage to trees, ranging from defoliation to snapped and uprooted trunks, and have been widely reported (Basnet et al., 1992; Brokaw & Gear, 1991; Curran, Gersbach, Edwards, & Krockenberger, 2008; Everham & Brokaw, 1996; Franklin, Drake, McConkey, Tonga, & Smith, 2004; Metcalfe et al., 2008; Turton, 2008; Zimmerman et al., 1994). The gaps and defoliation generate space and allow more light penetration on the forest floor, enhancing regeneration (e.g., Bellingham, Tanner, & Healey, 1995; Turner et al., 1998; Uriarte et al., 2004; Whitmore, 1974).

Because TCs tend to disproportionately affect big trees and encourage regeneration (Beard et al., 2005; Everham & Brokaw, 1996; Franklin, 2007; Hjerpe, Hedenas, & Elmqvist, 2001; Murphy, Metcalfe, Bradford, & Ford, 2014; Murphy et al., 2008; Roth, 1992), forests affected by frequent cyclones would be expected to have lower canopies and higher stem densities. Comparative regional studies support these expectations for Madagascar vs. African rainforests (De Gouvenain & Silander, 2003), New Caledonia vs. other Southwest Pacific humid forests (Ibanez et al., 2017), Caribbean vs. Neotropical dry forests (Quigley & Platt, 2003; Van Bloem, Murphy, & Lugo, 2007). Furthermore, stem density increased with TC frequency in lowland rainforests of five South Pacific archipelagos (Keppel et al., 2010), and was higher in Neotropical dry forests that experienced TCs in the last 25 years compared to regions with no TCs (Gillespie, Zutta, Early, & Saatchi, 2006). However, a pantropical test for these invisible, medium- to long-term effects (Lugo, 2008) on forest structure is still lacking.

The availability of high-resolution historical databases of TCs trajectories and intensities such as the International Best Track Archive for Climate Stewardship (IBTrACS; Knapp, Kruk, Levinson, Diamond, & Neumann, 2010) allows to objectively quantify the cyclone regime at any location (e.g., Schreck et al., 2014). The frequency or the intensity of TCs (i.e., the maximum sustained wind speed) is often used to estimate the level of disturbance induced by TCs in a given area. However, it has been suggested that this level of disturbance (or the destructiveness) is better described by the energy transferred by TCs to the land surface, as this parameter combines the frequency, intensity, size and translation speed of TCs (Camargo & Sobel, 2005; Emanuel, 2005).

In this study, we test on a pantropical scale the long-term, invisible effects (*sensu* Lugo, 2008) of TCs on the structure of tropical forests. We test it by analysing the effects of the frequency and intensity of TCs on structural features of forests (stem density, basal

area and canopy height), relative to other climatic variables (mean annual temperature [MAT] and water availability) and human disturbances (human foot print index) that are known to affect forest structure (e.g., Klein, Randin, & Körner, 2015; Moles et al., 2009; Tao, Guo, Li, Wang, & Fang, 2016; Zhang, Nielsen, Mao, Chen, & Svenning, 2016). We hypothesize that TC are an important disturbance that has long-term effects on the structure of tropical forests and that, therefore, areas exposed to frequent and intense TCs would exhibit higher stem densities and lower canopy heights. Our findings are highly relevant to understanding how forecasted changes in the behaviour of TCs (Christensen et al., 2013; Walsh et al., 2016) may impact forest ecosystems.

2 | MATERIALS AND METHODS

2.1 | Structural features

We compiled the location, stem density (number of stems per ha), basal area (the total cross section area in m²/ha, which is a good proxy of above-ground biomass; e.g., Slik et al., 2010), maximum size (maximum DBH, diameter at breast height, cm), mean canopy height (m) and maximum height for plants with stem diameter ≥10 cm at ~1.3 m above the base (DBH) from published inventories for plots ≥0.1 ha in size (Ibanez et al., 2018). Relevant literature was identified using key word searches in ISI Web of Science (<https://webof-knowledge.com/>) and Google Scholar (<http://scholar.google.com/>). Our final dataset was derived from 88 scientific papers published between 1983 and 2017. Montane forest plots (as defined by authors of source data, i.e., "pre-montane," "lower-montane," "montane" and "upper-montane") were not considered in the analysis (Aiba and Kitayama, 1999; Clark et al., 2015; Culmsee et al., 2011; Davidar, Mohandass, & Vijayan, 2007; Noumi, 2013; Rakotomalaza and Messmer, 1999; Yamada, 1975). We also only considered plots where authors did not report evidence of recent natural (e.g., TCs) or anthropogenic disturbances (e.g., removal of trees). When canopy height was reported with lower and higher bounds (e.g., 20–30 m), we used the mean value (e.g., 25 m).

2.2 | Cyclones

The frequency and energy released by TCs were computed over the 1981–2016 period from the IBTrACS database, an exhaustive and global database (<https://www.ncdc.noaa.gov/ibtracs/>, see Knapp et al., 2010). We assumed these values, derived from a 35-year period, to be representative of the relative TC regime in different regions over the Holocene on a global scale. Although some geomorphological records in TC-prone regions show centennial to millennial variations in local TC frequency and intensity during the Holocene, these variation are pseudo-cyclic and do not indicate long-term trends (e.g., Burn & Palmer, 2015; Haig, Nott, & Reichart, 2014; Nott & Forsyth, 2012; Toomey, Donnelly, & Woodruff, 2013). Instead, global patterns of TCs across different

regions vary less on a centennial scale but respond more strongly to major changes in global climate over longer time-scales, such as the Pliocene glaciations (e.g., Yoo, Galewsky, Camargo, Korty, & Zamora, 2016).

The IBTrACs database provides information on the position, maximum sustained wind speed and translation speed (an indicator of the duration of disturbance impact, as slower moving cyclones imply longer presence of damaging winds) of each inventoried cyclone at 6-hr intervals. We used these data to calculate the average frequency and energy released by TCs was computed as a proxy of the level of TC disturbance.

For each grid cell with a spatial resolution of 1° latitude \times 1° longitude ($\approx 12,345 \text{ km}^2$ at the equator, Figure 1), the energy released by TCs was computed as the power dissipated by friction in the surface layer (Emanuel, 2005) following Menkes et al. (2016) and Vincent et al. (2012). At each point, the maximum 10-min sustained wind speeds at 10 m above the ground (V) was extracted from the database. Only events reaching $V > 17 \text{ m/s}$ at some point were considered, i.e., tropical storms and category one to five TCs according to the Saffir–Simpson Hurricane Scale (Simpson, 1974). Following Vincent et al. (2012), the spatial pattern of each TC at each time step was computed using the Willoughby, Darling, and Rahn (2006) idealized vortex. We then computed the power dissipation index (PDI) as the integration over the entire lifetime (τ) and spatial extent (r_0) of TCs as follows:

$$\text{PDI} = 2\pi \int_0^\tau \int_0^{r_0} \alpha |V|^3 r dr$$

where α ($\alpha = C_D \rho$ with C_D being the surface drag coefficient and ρ the surface air density) is considered a constant (see Emanuel, 2005). The PDI was then integrated over each 1° latitude \times 1° longitude grid cells and averaged to get the average yearly PDI. As expected, TC frequency and PDI were strongly correlated (see Figure S1.1).

2.3 | Bioclimatic variables

Plots were classified as dry or humid forest plots following authors' classifications, with tropical dry forests being identified by the common occurrence of drought-deciduous canopy species. For each plot, we extracted the corresponding TC frequency and energy released by TCs (PDI). We also computed the MAT, mean annual precipitation (MAP), annual potential transpiration (PET) and a water availability index as MAP–PET. MAP and MAT were extracted from the WorldClim 2.0 database (<http://www.worldclim.org/version2>, Fick & Hijmans, 2017) and PET from the CGIAR–CSI database (<http://www.cgiar-csi.org/>, Trabucco & Zomer, 2009).

2.4 | Human disturbances

Human activities have had a major impact on forests and may affect forest structure directly, e.g., by cutting trees for wood (Crowther et al., 2015; Hansen et al., 2013), or indirectly, e.g., by edge effects

related to fragmentation (Laurance, 1997). To take into account the effects of these disturbances, we extracted the human foot print index (HFP, Human Footprint 2009, <http://wcshumanfootprint.org/>, Venter et al., 2016), as a proxy of potential human disturbances. This index measures the cumulative impact of direct pressures on natural ecosystems from human activities and includes the extent of built environments, crop land and pasture land, the human population density, the night-lights and the circulation ways (railways, roads and navigable waterways).

2.5 | Analysis

All analyses were performed using R.3.3.0 (R Core Team, 2016). All explanatory variables (except TC frequency and PDI) were extracted on 10-min ($\sim 340 \text{ km}^2$) spatial resolution maps. To avoid pseudo-replication, we averaged structural and climatic features within 1×1 degree grid cells (i.e., the spatial resolution of the PDI maps) and forest type (dry and humid forests). We used a total of 438 plots (355 in humid forests, 83 in dry forests) that were pooled into 250 grid cells (203 in humid forests, 47 in dry forests, Figure 1 and Figure S1.2). Multicollinearity was <0.5 (Spearman's rho) for all explanatory variables (Table S1.1).

We first tested whether structural features differed between areas experiencing low- (freq. <0.01 TC/year, i.e., less than one TC per century is expected based on the 35-year IBTrACs dataset), medium- ($0.01 \geq \text{freq.} < 0.1$ TC/year, i.e., 1–10 TCs per century) or high- (freq. ≥ 0.1 TC/year, i.e., at least one TC per decade) TC frequency. This was done separately for dry and humid forests (see distribution of TC frequency Figure S1.1). The significance of the differences was tested using one-way pairwise Wilcoxon rank sum tests with corrections for multiple testing (Holm's correction).

We used multivariate linear regressions to test the relative importance of PDI and other relevant parameters; MAT, water availability (MAP–PET) and human disturbances (HFP). Response variables describing the structure of forests were log-transformed to increase normality (Figure S1.3). Because basal area and maximum size, and canopy height and maximum height, were highly linearly correlated for both forest types, we focussed on three relatively independent structural variables: stem density, basal area and canopy height (Figure S1.4). We used the *MuMIn* R package (Bartoń, 2016) and the *dredge* function to generate different sets of models representing all possible combinations and subsets of explanatory variables. We then selected the best models based on the Akaike information criterion ($\Delta\text{AIC} < 2$ from the best models, Burnefeld & Phillimore, 2012). Explanatory variables were centred and scaled before fitting models to allow a fair comparison of their respective effects.

3 | RESULTS

Across the tropics and in both forest types, forests located in areas experiencing high TC frequency (>1 per decade) exhibited higher




FIGURE 2 Comparison of structural features in dry and humid forests with different tropical cyclone (TC) frequencies (<0.01 = low frequency, less than one TC per century, 0.01–0.1 = medium frequency, at least one TC per century and >0.1 = high frequency, at least one TC per decade). Letters above groups indicate significant differences, i.e., forest classes sharing a letter are not significantly different ($p > 0.05$) using a pairwise Wilcoxon rank sum test

stem densities (Figure 2). The stem density in these regions was on average 798 ± 253 stems per ha in dry forests and 888 ± 340 stems per ha in humid forests, compared to 643 ± 195 stems per ha and 703 ± 194 stems per ha, respectively, in regions experiencing less than one TC per decade. In contrast, basal area did not differ significantly between the three cyclone frequency classes (low, medium and high) for both dry and humid forests. The canopy of humid forests (27 ± 8 m) was on average more than twice as high as in dry forests (12 ± 5 m). Humid forests exposed to frequent TCs were also significantly shorter (20 ± 6 m) than those exposed to low- and medium-TC frequencies (27 ± 8 m).

The amount of energy released by TCs (the so-called, PDI) was positively correlated to stem density and basal area (Table 1). PDI together with MAT explained 14% of stem density variance, but had no significant effect basal area (Table 1 and Figure 3). Canopy height had a significant positive relationship with water availability and a significant negative relationship with human foot print. Water availability was significantly lower (-177.7 ± 626.65 mm vs. $700.6 \pm 1,030.1$ mm; Wilcoxon rank test $p < 0.001$, Figure S1.4) and the human footprint significantly higher (12.1 ± 9.1 vs. 8.0 ± 6.4 ; Wilcoxon rank test $p < 0.01$, Figure S1.5) in dry forests compared to humid forests.

TABLE 1 Best multivariate linear regressions explaining structural features in tropical forests overall and in humid and dry forest separately

Forest types	Structural features	N	Model ranks	Coefficient estimate and significance				Model performance	
				PDI	MAT	MAP-PET	HFP	R ²	AICc
Both	Stem density	245	1	0.10***	−0.08***			0.14	119.24
			2	0.10***	−0.08***		0.01	0.14	121.24
	Basal area	229	1		−0.09***	0.05 [†]		0.06	191.88
			2		−0.08**			0.05	192.92
			3		−0.08**	0.05	−0.01	0.06	193.44
			4	0.01	−0.08**	0.05 [†]	−0.05 [†]	0.06	193.77
	Canopy height	78	1		0.12*	0.19***	−0.20***	0.32	93.08
			2	−0.06	0.10	0.19***	−0.18**	0.33	94.02
			3	−0.09 [†]		0.18***	−0.16**	0.30	94.52
Humid	Stem density	201	1	0.08**	−0.09***			0.17	74.03
			2	0.08**	−0.09***		0.02	0.17	75.50
	Basal area	190	1		−0.09***			0.08	119.41
			2	0.04	−0.09***			0.09	119.63
			3		−0.09***		0.03	0.08	120.68
			4	0.04	−0.09***		0.02	0.09	121.17
	Canopy height	51	1	−0.15***		0.07 [†]		0.33	23.42
			2	−0.13**	0.06	0.07 [†]		0.36	24.05
			3	−0.15***				0.29	24.10
			4	−0.13**	0.06			0.31	24.90
Dry	Stem density	44	1	0.13*				0.11	39.65
			2	0.11 [†]		0.13		0.15	40.64
			3			0.18		0.07	41.28
	Basal area	39	1		0.18 [†]		−0.15*	0.17	44.39
			2						45.13
			3				−0.10	0.07	45.38
			4		0.10			0.04	46.32
	Canopy height ^a	27	1			−0.22		0.12	25.62
			2						25.79
			3				−0.09	0.10	26.10
			4			−0.20	−0.08	0.19	26.51

Note. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$; [†] $p < 0.1$.

^aOnly the four models with the lowest AICc are shown.

PDI, power dissipation index; MAT, mean annual temperature; MAP-PET, mean annual precipitations – potential evapotranspiration; HFP, human foot print.

Within humid forests, which was less affected by human disturbance than dry forest, PDI was a significant predictor of stem density and canopy height (Table 1), explaining alone about 10% and 29% of total variation in stem density and canopy height respectively (Figure 3). MAT was the other significant predictor. Together with PDI it explained 17% and 9% of total variance of stem density and basal area (Table 1). PDI had positive relationships with stem density, while MAT had positive relationships with stem density and basal area. Canopy height tended to increase with water availability which explained together with PDI 33% of the observed variance. Within

dry forest, PDI was the only significant predictor of stem density, explaining about 11% of its variance (Figure 3). HFP together with MAT explained about 17% of the variance in basal area, while none of the variables was significantly related to canopy height variations.

4 | DISCUSSION

Our results confirm that TCs are important disturbances that have a long-term, global-scale effects on forest structure, despite being




FIGURE 3 Relationships between the energy released by tropical cyclones (power dissipation index), stem density and canopy height (on log scales). Thick lines represent significant linear relationships (** $p < 0.001$, * $p < 0.05$)

relatively uncommon weather events (about 80–90 TCs per year, see Schreck et al., 2014; Maue, 2011). We show for the first time that stem density and canopy height are significantly related to the frequency of and energy released by TCs in a globally consistent way. Furthermore, our results indicate that these relationships were most pronounced for stem density and for humid forests, which are not limited by water availability and are less impacted by human disturbances (based on lower human foot print) than dry forests. Therefore, forecasted changes in the intensities, sizes and trajectories of TCs are likely to have important impacts on the structure of tropical and subtropical forests (Kossin, Emanuel, & Vecchi, 2014; Mei, Xie, Primeau, McWilliams, & Pasquero, 2015).

Our results explained a considerable proportion of the variation in stem density and canopy height in our dataset, suggesting that TCs are important large-scale disturbances that have long-term impacts on forests structure. Our analysis did not capture edaphic and topographic variation known to impact forest structure from local to regional scales, which makes the strong relationship we found between cyclone and forest structure even more remarkable. Indeed, soil depth, nutrient availability and topographical position are known to affect forest structure (e.g., Clark & Clark, 2000; Quesada et al., 2012; Schut et al., 2014; Slik et al., 2010; Webb, Stanfield, & Jensen, 1999).

The strong, universal relationship between TCs and stem density documented in this study is not surprising. TCs are disturbances

that reduce the canopy cover of forests (Comita et al., 2009; Grove, Turton, & Siegenthaler, 2000; Hjerpe et al., 2001; Staben & Evans, 2008) and cause tree mortality (Webb et al., 2011; Zeng et al., 2009; Zimmerman et al., 1994). This results in increased light availability for several years in the undergrowth (Bellingham, Tanner, Rich, & Goodland, 1996; Lin et al., 2011; Luke, McLaren, & Wilson, 2014; Turton, 1992), creating suitable conditions for regeneration and the release of saplings (Luke et al., 2014; Nicotra, Chazdon, & Iriarte, 1999; Tanner, Rodriguez-Sanchez, Healey, Holdaway, & Bellingham, 2014). Therefore, frequent cyclones (e.g., one per decade) would be expected to maintain high light availability and, consequently, regeneration. Furthermore, higher mortalities and regeneration reported from TC-prone forests suggest that such forests may experience higher turnover and quicker dynamics than forests not or rarely affected by these disturbances (Swenson et al., 2012; Webb et al., 2011).

Given that TCs are known to trim forest canopies to lower heights (Brokaw & Gear, 1991; Lin et al., 2011), the moderate effect of TCs on canopy height may seem surprising. Our results suggest that canopy height first depends on hydraulic limitation and then by mechanical pruning by intense TC winds, as canopy height of dry forest (with MAP-PET being on average -177.7 ± 626.7 mm) was less than half that of humid forest (with MAP-PET being on average $700.6 \pm 1,030.1$ mm). This is in the line with recent global analyses showing canopy height to increase with water availability (Klein

et al., 2015; Zhang et al., 2016) and the hydraulic limitation hypothesis (Ryan, Phillips, & Bond, 2006; Ryan & Yoder, 1997).

In humid forests, frequent TC disturbances may result in a greater investment in horizontal over vertical growth, as suggested by the negative relationship between the energy released by TCs with canopy height and its weaker, positive relationship with basal area. This deduction is supported by recent studies comparing height-diameter tree allometries across the tropics, which have shown that areas exposed to frequent strong winds, such as New Caledonia (Blanchard et al., 2016) or Dominica (Thomas, Martin, & Mycroft, 2015), have lower growth in height relative to diameter. These changes in tree allometry could result from repeated pruning of crown tips by TCs (Brokaw & Gear, 1991) or from evolutionary adaptations to TCs, which have been suggested to explain differential responses of neotropical palms (Areaceae) to TCs (Griffith, Noblick, Dowe, Husby, & Calonje, 2008). The latter would imply TCs to have important evolutionary influences. Alternatively lower canopy heights may be due to resprouting, which is commonly observed in trees surviving TC damages (Bellingham, Tanner, & Healey, 1994; Scanlon, Petit, Tuiwawa, & Naikatini, 2018; Van Bloem et al., 2007; Zimmerman et al., 1994) and produces multitemmed trees that tend to be lower in height than single-stemmed trees (Givnish, 1984; Kruger, Midgley, & Cowling, 1997).

The expected changes in the frequency, intensity or geographical distribution of TCs due to anthropogenic climate change are therefore likely to impact the structure and dynamics of affected forests. While the influence of global warming on TCs remains uncertain (Walsh et al., 2016), there is an emerging consensus that TCs will increase in intensity (Christensen et al., 2013; Kang & Elsner, 2015; Mei et al., 2015), attain unprecedented sizes and intensities (Lin & Emanuel, 2016; Mei et al., 2015), and expand their trajectories poleward (Kossin et al., 2014). Our findings suggest that more intense TCs may alter tropical forest structure (density, height and basal area) and dynamics (higher turnover). Similar changes can be expected in forests at higher latitudes experiencing TCs for the first time. Furthermore, forests subjected to more intense cyclones may experience changes in species composition (Keppel et al., 2010; Webb et al., 2011) and increased vulnerability to drought and other threats, such as fire and invasion by weeds (Beard et al., 2005; Franklin, 2007; Hjerpe et al., 2001; Murphy et al., 2008).

The significant, negative relationship of human foot print with forest canopy height and basal area highlights that factors other than TC are impacting forest structure. This relationship could be the result of selective tree removal by humans or edge effects related to fragmentation (Laurance, Delamonica, Laurance, Vasconcelos, & Lovejoy, 2000; Lindenmayer, Laurance, & Franklin, 2012). While these potential causes are purely speculative, it is well documented that forests are being degraded and destroyed at alarming rates (Crowther et al., 2015; Hansen et al., 2013). A direct anthropogenic influence on forest structure (and hence dynamics) is further supported by the effects of the human foot print being stronger in tropical dry forests, which is known to be extremely disturbed and fragmented (Gillespie et al., 2014; Janzen, 1988; Miles et al., 2006).

5 | CONCLUSIONS

We provide the first evidence that the visible, short-term damages caused by TCs have less obvious, long-term effects on the structure of forests on a pantropical scale. The consistent finding of an increased stem density in regions experiencing a stronger TC disturbance regime suggests that TCs alter the structure of forests in a globally consistent manner. Furthermore, our results suggest that, within humid forests, TCs may have a stronger effect on canopy height than water availability. This implies that the predicted higher intensities and higher latitudes of cyclone impact in future will result in major and predictable changes in forest structure and, hence, global carbon stocks.

ACKNOWLEDGEMENTS

T.I. was funded by the Agency for Economic and Environmental Development (DDEE) of the North Province of New Caledonia and G.K. was supported by an Alexander von Humboldt fellowship. We thank all people who contributed to collecting plot data. We are notably most grateful to the late Alwyn H. Gentry, the Missouri Botanical Garden, and collectors who assisted Gentry in several studied islands. We thank two anonymous reviewers and Peter Bellingham for helpful comments on the manuscript.

AUTHORS' CONTRIBUTIONS

T.I., G.K. and P.B. conceived of the idea; T.I., G.K., T.W.G., G.R. and P.B. compiled the forest inventory data; C.M. and M.L. computed the tropical cyclone maps; T.I., G.K. and M.M. analysed the data; T.I. and G.K. interpreted the results and drafted the paper. All authors contributed to the writing of the paper.

DATA ACCESSIBILITY

Data available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.4nn33b2> (Ibanez et al., 2018).

ORCID

Thomas Ibanez  <http://orcid.org/0000-0002-3192-1721>

Gunnar Keppel  <http://orcid.org/0000-0001-7092-6149>

Gonzalo Rivas-Torres  <http://orcid.org/0000-0002-2704-8288>

REFERENCES

- Aiba, S., & Kitayama, K. (1999). Structure, composition and species diversity in an altitude-substrate matrix of rain forest tree communities on Mount Kinabalu, Borneo. *Plant Ecology*, 140, 139–157. <https://doi.org/10.1023/A:1009710618040>
- Bartoň, K. (2016) {MuMIn}: multi-model inference, {R} package version 1.15.6. Retrieved from <http://cran.r-project.org/package=MuMIn>
- Basnet, K., Likens, G. E., Scatena, F. N., & Lugo, A. E. (1992). Hurricane Hugo: Damage to a tropical rain forest in Puerto Rico. *Journal of Tropical Ecology*, 8, 47–55. <https://doi.org/10.1017/S0266467400006076>

- Beard, K. H., Vogt, K. A., Vogt, D. J., Scatena, F. N., Covich, A. P., Sigurdardottir, R., ... Crowl, T. A. (2005). Structural and functional responses of a subtropical forest to 10 years of hurricanes and droughts. *Ecological Monographs*, 75, 345–361. <https://doi.org/10.1890/04-1114>
- Bellingham, P. J., Tanner, E. V. J., & Healey, J. R. (1994). Sprouting of trees in Jamaican montane forests, after a hurricane. *Journal of Ecology*, 82, 747–758. <https://doi.org/10.2307/2261440>
- Bellingham, P. J., Tanner, E. V. J., & Healey, J. R. (1995). Damage and responsiveness of Jamaican montane tree species after disturbance by a hurricane. *Ecology*, 76, 2562–2580. <https://doi.org/10.2307/2265828>
- Bellingham, P. J., Tanner, E. V. J., Rich, P. M., & Goodland, T. C. R. (1996). Changes in light below the canopy of a Jamaican montane rainforest after a hurricane. *Journal of Tropical Ecology*, 12, 699–722. <https://doi.org/10.1017/S0266467400009883>
- Blanchard, E., Birnbaum, P., Ibanez, T., Boutreux, T., Antin, C., Ploton, P., ... Couteron, P. (2016). Contrasted allometries between stem diameter, crown area, and tree height in five tropical biogeographic areas. *Trees*, 30, 1953–1968.
- Boose, E. R., Serrano, M. I., & Foster, D. R. (2004). Landscape and regional impacts of hurricanes in Puerto Rico. *Ecological Monographs*, 74, 335–352. <https://doi.org/10.1890/02-4057>
- Brokaw, N. V. L., & Grear, J. S. (1991). Forest structure before and after hurricane Hugo at 3 elevation in the Luguillo mountains, Puerto Rico. *Biotropica*, 23, 386–392. <https://doi.org/10.2307/2388256>
- Bunnefeld, N., & Phillimore, A. B. (2012). Island, archipelago and taxon effects: Mixed models as a means of dealing with the imperfect design of nature's experiments. *Ecography*, 35, 15–22. <https://doi.org/10.1111/j.1600-0587.2011.07078.x>
- Burn, M. J., & Palmer, S. E. (2015). Atlantic hurricane activity during the last millennium. *Scientific Reports*, 5, 12838. <https://doi.org/10.1038/srep12838>
- Roth, L. C. (1992). Hurricanes and mangrove regeneration: Effects of hurricane Joan, October 1988, on the vegetation of Isla del Venado, Bluefields, Nicaragua. *Biotropica*, 24, 375–384. <https://doi.org/10.2307/2388607>
- Camargo, S. J., & Sobel, A. H. (2005). Western North Pacific tropical cyclone intensity and ENSO. *Journal of Climate*, 18, 2996–3006. <https://doi.org/10.1175/JCLI3457.1>
- Chan, K. T. F., & Chan, J. C. L. (2015). Global climatology of tropical cyclone size as inferred from QuikSCAT data. *International Journal of Climatology*, 35, 4843–4848. <https://doi.org/10.1002/joc.4307>
- Christensen, J. H., Krishna Kumar, K., Aldrian, E., An, S.-I., Cavalcanti, I. F. A., deCastro, M., & Zhou, T. (2013). Climate phenomena and their relevance for future regional climate change. In T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P. M. Midgley (Eds.), *Climate change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 92). New York, NY: Cambridge University Press.
- Clark, D. B., & Clark, D. A. (2000). Landscape-scale variation in forest structure and biomass in a tropical rain forest. *Forest Ecology and Management*, 137, 185–198. [https://doi.org/10.1016/S0378-1127\(99\)00327-8](https://doi.org/10.1016/S0378-1127(99)00327-8)
- Clark, D. B., Hurtado, J., & Saatchi, S. S. (2015). Tropical rain forest structure, tree growth and dynamics along a 2700-m elevational transect in Costa Rica. *PLoS ONE*, 10, e0122905. <https://doi.org/10.1371/journal.pone.0122905>
- Comita, L. S., Uriarte, M., Thompson, J., Jonckheere, I., Canham, C. D., & Zimmerman, J. K. (2009). Abiotic and biotic drivers of seedling survival in a hurricane-impacted tropical forest. *Journal of Ecology*, 97, 1346–1359. <https://doi.org/10.1111/j.1365-2745.2009.01551.x>
- Costanza, R., Pérez-Maqueo, O., Martínez, M. L., Sutton, P., Anderson, S. J., & Mulder, K. (2008). The value of coastal wetlands for hurricane protection. *Ambio*, 37, 241–248. [https://doi.org/10.1579/0044-7447\(2008\)37\[241:TVOCWF\]2.0.CO;2](https://doi.org/10.1579/0044-7447(2008)37[241:TVOCWF]2.0.CO;2)
- Crowther, T. W., Glick, H. B., Covey, K. R., Bettigole, C., Maynard, D. S., Thomas, S. M., ... Bradford, M. A. (2015). Mapping tree density at a global scale. *Nature*, 525, 201–205. <https://doi.org/10.1038/nature14967>
- Culmsee, H., Pitopang, R., Mangopo, H., & Sabir, S. (2011). Tree diversity and phytogeographical patterns of tropical high mountain rain forests in Central Sulawesi, Indonesia. *Biodiversity and Conservation*, 20, 1103–1123. <https://doi.org/10.1007/s10531-011-0019-y>
- Curran, T. J., Gersbach, L. N., Edwards, W., & Krockenberger, A. K. (2008). Wood density predicts plant damage and vegetative recovery rates caused by cyclone disturbance in tropical rainforest tree species of North Queensland, Australia. *Austral Ecology*, 33, 442–450. <https://doi.org/10.1111/j.1442-9993.2008.01899.x>
- Davidar, P., Mohandass, D., & Vijayan, S. L. (2007). Floristic inventory of woody plants in a tropical montane (shola) forest in the Palni hills of the Western Ghats, India. *Tropical Ecology*, 48, 15–25.
- De Gouvenain, R. C., & Silander, J. A. (2003). Do tropical storm regimes influence the structure of tropical lowland rain forests? *Biotropica*, 35, 166–180.
- Emanuel, K. (2005). Increasing destructiveness of tropical cyclones over the past 30 years. *Nature*, 436, 686–688. <https://doi.org/10.1038/nature03906>
- Everham, E. M., & Brokaw, N. V. L. (1996). Forest damage and recovery from catastrophic wind. *Botanical Review*, 62, 113–185. <https://doi.org/10.1007/BF02857920>
- Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, 37, 4302–4315. <https://doi.org/10.1002/joc.5086>
- Franklin, J. (2007). Recovery from clearing, cyclone and fire in rain forests of Tonga, South Pacific: Vegetation dynamics 1995–2005. *Austral Ecology*, 32, 789–797. <https://doi.org/10.1111/j.1442-9993.2007.01766.x>
- Franklin, J., Drake, D. R., McConkey, K. R., Tonga, F., & Smith, L. B. (2004). The effects of Cyclone Waka on the structure of lowland tropical rain forest in Vava'u, Tonga. *Journal of Tropical Ecology*, 20, 409–420. <https://doi.org/10.1017/S0266467404001543>
- Gillespie, T. W., O'Neill, K., Keppel, G., Pau, S., Meyer, J.-Y., Price, J. P., & Jaffré, T. (2014). Prioritizing conservation of tropical dry forests in the Pacific. *Oryx*, 48, 337–344. <https://doi.org/10.1017/S0030605313000264>
- Gillespie, T. W., Zutta, B. R., Early, M. K., & Saatchi, S. (2006). Predicting and quantifying the structure of tropical dry forests in South Florida and the Neotropics using spaceborne imagery. *Global Ecology and Biogeography*, 15, 225–236. <https://doi.org/10.1111/j.1466-822X.2005.00203.x>
- Givnish, T. J. (1984). Leaf and canopy adaptations in tropical forests. In E. Medina, H. A. Mooney & C. Vázquez-yánes (Eds.), *Physiological ecology of plants of the wet tropics* (pp. 51–84). The Hague, the Netherlands: Dr W. Junk Publishers. <https://doi.org/10.1007/978-94-009-7299-5>
- Griffith, M. P., Noblick, L. R., Dowe, J. L., Husby, C. E., & Calonje, M. A. (2008). Cyclone tolerance in new world arecaceae: Biogeographic variation and abiotic natural selection. *Annals of Botany*, 102, 591–598. <https://doi.org/10.1093/aob/mcn132>
- Grove, S. J., Turton, S. M., & Siegenthaler, D. T. (2000). Mosaics of canopy openness induced by tropical cyclones in lowland rain forests with contrasting management histories in northeastern Australia. *Journal of Tropical Ecology*, 16, 883–894. <https://doi.org/10.1017/S0266467400001784>
- Haig, J., Nott, J., & Reichart, G.-J. (2014). Australian tropical cyclone activity lower than at any time over the past 550–1,500 years. *Nature*, 505, 667. <https://doi.org/10.1038/nature12882>
- Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., ... Townshend, J. R. G. (2013). High-resolution global maps of 21st-century forest cover change. *Science*, 342, 850–853. <https://doi.org/10.1126/science.1244693>

- Hjerpe, J., Hedenas, H., & Elmqvist, T. (2001). Tropical rain forest recovery from cyclone damage and fire in Samoa. *Biotropica*, 33, 249–259. <https://doi.org/10.1111/j.1744-7429.2001.tb00176.x>
- Ibanez, T., Blanchard, E., Hequet, V., Keppel, G., Laidlaw, M., Pouteau, R., ... Birnbaum, P. (2017). High endemism and stem density distinguish New Caledonian from other high-diversity rainforests in the Southwest Pacific. *Annals of Botany*, 121, 25–35.
- Ibanez, T., Keppel, G., Menkes, C., Gillespie, T. W., Lengaigine, M., Mangeas, M., ... Birnbaum, P. (2018). Data from: Globally consistent impact of tropical cyclones on the structure of tropical and subtropical forests. *Dryad Digital Repository*, <https://doi.org/10.5061/dryad.4nn33b2>
- Janzen, D. H. (1988). Tropical dry forests – The most endangered major tropical ecosystem. In E. O. Wilson & F. M. Peter (Eds.), *Biodiversity* (pp. 130–137). Washington, DC: National Academies Press.
- Kang, N.-Y., & Elsner, J. B. (2015). Trade-off between intensity and frequency of global tropical cyclones. *Nature Climate Change*, 5, 661–664. <https://doi.org/10.1038/nclimate2646>
- Keppel, G., Buckley, Y. M., & Possingham, H. P. (2010). Drivers of lowland rain forest community assembly, species diversity and forest structure on islands in the tropical South Pacific. *Journal of Ecology*, 98, 87–95. <https://doi.org/10.1111/j.1365-2745.2009.01595.x>
- Klein, T., Randin, C., & Körner, C. (2015). Water availability predicts forest canopy height at the global scale. *Ecology Letters*, 18, 1311–1320. <https://doi.org/10.1111/ele.12525>
- Knaff, J. A., Longmore, S. P., & Molenaar, D. A. (2014). An objective satellite-based tropical cyclone size climatology. *Journal of Climate*, 27, 455–476. <https://doi.org/10.1175/JCLI-D-13-00096.1>
- Knapp, K. R., Kruk, M. C., Levinson, D. H., Diamond, H. J., & Neumann, C. J. (2010). The international best track archive for climate stewardship (IBTrACS). *Bulletin of the American Meteorological Society*, 91, 363–376. <https://doi.org/10.1175/2009BAMS2755.1>
- Kossin, J. P., Emanuel, K. A., & Vecchi, G. A. (2014). The poleward migration of the location of tropical cyclone maximum intensity. *Nature*, 509, 349–352. <https://doi.org/10.1038/nature13278>
- Kruger, L. M., Midgley, J. J., & Cowling, R. M. (1997). Resprouters vs reseeder in South African forest trees; a model based on forest canopy height. *Functional Ecology*, 11, 101–105. <https://doi.org/10.1046/j.1365-2435.1997.00064.x>
- Laurance, W. F. (1997). Hyper-disturbed parks: Edge effects and the ecology of isolated rainforest reserves in tropical Australia. In W. F. Laurance, & R. O. Bierregaard (Eds.), *Tropical forest remnants: Ecology, Management, and conservation of fragmented communities* (pp. 71–84). Chicago, IL: The University of Chicago Press.
- Laurance, W. F., Delamonica, P., Laurance, S. G., Vasconcelos, H. L., & Lovejoy, T. E. (2000). Conservation: Rainforest fragmentation kills big trees. *Nature*, 404, 836–836. <https://doi.org/10.1038/35009032>
- Le Page, M. (2016). Fiji storm breaks records. *New Scientist*, 229, 6. [https://doi.org/10.1016/S0262-4079\(16\)32303-X](https://doi.org/10.1016/S0262-4079(16)32303-X)
- Lin, N., & Emanuel, K. (2016). Grey swan tropical cyclones. *Nature Climate Change*, 6, 106–111. <https://doi.org/10.1038/nclimate2777>
- Lin, T.-C., Hamburg, S. P., Lin, K.-C., Wang, L.-J., Chang, C.-T., Hsia, Y.-J., ... Liu, C.-P. (2011). Typhoon disturbance and forest dynamics: Lessons from a Northwest Pacific subtropical forest. *Ecosystems*, 14, 127–143. <https://doi.org/10.1007/s10021-010-9399-1>
- Lindenmayer, D. B., Laurance, W. F., & Franklin, J. F. (2012). Global decline in large old trees. *Science*, 338, 1305–1306. <https://doi.org/10.1126/science.1231070>
- Lugo, A. E. (2008). Visible and invisible effects of hurricanes on forest ecosystems: An international review. *Austral Ecology*, 33, 368–398. <https://doi.org/10.1111/j.1442-9993.2008.01894.x>
- Luke, D., McLaren, K., & Wilson, B. (2014). The effects of a hurricane on seedling dynamics and abiotic interactions in a tropical lower montane wet forest. *Journal of Tropical Ecology*, 30, 55–66. <https://doi.org/10.1017/S0266467413000734>
- Maue, R. N. (2011). Recent historically low global tropical cyclone activity. *Geophysical Research Letters*, 38, L14803.
- Mei, W., Xie, S.-P., Primeau, F., McWilliams, J. C., & Pasquero, C. (2015). Northwestern Pacific typhoon intensity controlled by changes in ocean temperatures. *Science Advances*, 1, e1500014.
- Menkes, C. E., Lengaigine, M., Lévy, M., Ethé, C., Bopp, L., Aumont, O., ... Jullien, S. (2016). Global impact of tropical cyclones on primary production. *Global Biogeochemical Cycles*, 30, 767–786. <https://doi.org/10.1002/2015GB005214>
- Metcalf, D. J., Bradford, M. G., & Ford, A. J. (2008). Cyclone damage to tropical rain forests: Species- and community-level impacts. *Austral Ecology*, 33, 432–441. <https://doi.org/10.1111/j.1442-9993.2008.01898.x>
- Miles, L., Newton, A. C., DeFries, R. S., Ravilious, C., May, I., Blyth, S., ... Gordon, J. E. (2006). A global overview of the conservation status of tropical dry forests. *Journal of Biogeography*, 33, 491–505. <https://doi.org/10.1111/j.1365-2699.2005.01424.x>
- Moles, A. T., Warton, D. I., Warman, L., Swenson, N. G., Laffan, S. W., Zanne, A. E., ... Leishman, M. R. (2009). Global patterns in plant height. *Journal of Ecology*, 97, 923–932. <https://doi.org/10.1111/j.1365-2745.2009.01526.x>
- Murphy, H. T., Metcalfe, D. J., Bradford, M. G., Ford, A. F., Galway, K. E., Sydes, T. A., & Westcott, D. J. (2008). Recruitment dynamics of invasive species in rainforest habitats following Cyclone Larry. *Austral Ecology*, 33, 495–502. <https://doi.org/10.1111/j.1442-9993.2008.01904.x>
- Murphy, H. T., Metcalfe, D. J., Bradford, M. G., & Ford, A. J. (2014). Community divergence in a tropical forest following a severe cyclone. *Austral Ecology*, 39, 696–709. <https://doi.org/10.1111/aec.12133>
- Nicotra, A. B., Chazdon, R. L., & Iriarte, S. V. B. (1999). Spatial heterogeneity of light and woody seedling regeneration in tropical wet forests. *Ecology*, 80, 1908–1926. [https://doi.org/10.1890/0012-9658\(1999\)080\[1908:SHOLAW\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[1908:SHOLAW]2.0.CO;2)
- Nott, J., & Forsyth, A. (2012). Punctuated global tropical cyclone activity over the past 5,000 years. *Geophysical Research Letters*, 39, L14703.
- Noumi, E. (2013). Floristic inventory of woody species in the Manengouba Mountain forest, Cameroon. *Journal of Biology and Life Science*, 4, 28.
- Quesada, C. A., Phillips, O. L., Schwarz, M., Czimczik, C. I., Baker, T. R., Patiño, S., ... Lloyd, J. (2012). Basin-wide variations in Amazon forest structure and function are mediated by both soils and climate. *Biogeosciences*, 9, 2203–2246. <https://doi.org/10.5194/bg-9-2203-2012>
- Quigley, M. F., & Platt, W. J. (2003). Composition and structure of seasonally deciduous forests in the Americas. *Ecological Monographs*, 73, 87–106. [https://doi.org/10.1890/0012-9615\(2003\)073\[0087:CASOSD\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2003)073[0087:CASOSD]2.0.CO;2)
- R Core Team. (2016). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Rakotomalaza, P. J., & Messmer, N. (1999). Structure and floristic composition of the vegetation in the Réserve Naturelle Intégrale d'Andohahela, Madagascar. In S. M. Goodman (Ed.), *A floral and faunal inventory of the Réserve Naturelle Intégrale d'Andohahela, Madagascar: With reference to elevational variation* (pp. 51–96). Chicago, IL: Field Museum of Natural History.
- Ryan, M. G., Phillips, N., & Bond, B. J. (2006). The hydraulic limitation hypothesis revisited. *Plant Cell and Environment*, 29, 367–381. <https://doi.org/10.1111/j.1365-3040.2005.01478.x>
- Ryan, M. G., & Yoder, B. J. (1997). Hydraulic limits to tree height and tree growth. *BioScience*, 47, 235–242. <https://doi.org/10.2307/1313077>
- Scanlon, A. T., Petit, S., Tuiwawa, M., & Naikati, A. (2018). Response of primary and secondary rainforest flowers and fruits to a cyclone, and implications for plant-servicing bats. *Global Change Biology*, 24, 3820–3836. <https://doi.org/10.1111/gcb.14103>
- Schreck, C. J. III, Knapp, K. R., & Kossin, J. P. (2014). The impact of best track discrepancies on global tropical cyclone climatologies using IBTrACS. *Monthly Weather Review*, 142, 3881–3899. <https://doi.org/10.1175/MWR-D-14-00021.1>

- Schut, A. G. T., Wardell-Johnson, G. W., Yates, C. J., Keppel, G., Baran, I., Franklin, S. E., Byrne, M. (2014). Rapid characterisation of vegetation structure to predict refugia and climate change impacts across a global biodiversity hotspot. *PLoS ONE*, 9, e82778. <https://doi.org/10.1371/journal.pone.0082778>
- Simpson, R. H. (1974). The hurricane disaster potential scale. *Weatherwise*, 27, 169–186.
- Slik, J. W. F., Aiba, S.-I., Brearley, F. Q., Cannon, C. H., Forshed, O., Kitayama, K., ... van Valkenburg, J. L. C. H. (2010). Environmental correlates of tree biomass, basal area, wood specific gravity and stem density gradients in Borneo's tropical forests. *Global Ecology and Biogeography*, 19, 50–60. <https://doi.org/10.1111/j.1466-8238.2009.00489.x>
- Staben, G. W., & Evans, K. G. (2008). Estimates of tree canopy loss as a result of Cyclone Monica, in the Magela Creek catchment northern Australia. *Austral Ecology*, 33, 562–569. <https://doi.org/10.1111/j.1442-9993.2008.01911.x>
- Swenson, N. G., Stegen, J. C., Davies, S. J., Erickson, D. L., Forero-Montaña, J., Hurlbert, A. H., ... Zimmerman, J. K. (2012). Temporal turnover in the composition of tropical tree communities: Functional determinism and phylogenetic stochasticity. *Ecology*, 93, 490–499. <https://doi.org/10.1890/11-1180.1>
- Tanner, E. V. J., Rodriguez-Sanchez, F., Healey, J. R., Holdaway, R. J., & Bellingham, P. J. (2014). Long-term hurricane damage effects on tropical forest tree growth and mortality. *Ecology*, 95, 2974–2983. <https://doi.org/10.1890/13-1801.1>
- Tao, S., Guo, Q., Li, C., Wang, Z., & Fang, J. (2016). Global patterns and determinants of forest canopy height. *Ecology*, 97, 3265–3270. <https://doi.org/10.1002/ecy.1580>
- Thomas, S. C., Martin, A. R., & Mycroft, E. E. (2015). Tropical trees in a wind-exposed island ecosystem: Height-diameter allometry and size at onset of maturity. *Journal of Ecology*, 103, 594–605. <https://doi.org/10.1111/1365-2745.12378>
- Toomey, M. R., Donnelly, J. P., & Woodruff, J. D. (2013). Reconstructing mid-late Holocene cyclone variability in the Central Pacific using sedimentary records from Tahaa, French Polynesia. *Quaternary Science Reviews*, 77, 181–189. <https://doi.org/10.1016/j.quascirev.2013.07.019>
- Trabucco, A., & Zomer, R. J. (2009). Global aridity index (global-aridity) and global potential evapo-transpiration (global-PET) geospatial database. Published online, available from the CGIAR-CSI GeoPortal at: <http://www.csi.cgiar.org>
- Turner, M. G., Baker, W. L., Peterson, C. J., & Peet, R. K. (1998). Factors influencing succession: Lessons from large, infrequent natural disturbances. *Ecosystems*, 1, 511–523. <https://doi.org/10.1007/s100219900047>
- Turner, M. G., & Dale, V. H. (1998). Comparing large, infrequent disturbances: What have we learned? *Ecosystems*, 1, 493–496. <https://doi.org/10.1007/s100219900045>
- Turner, M. G., Dale, V. H., & Everham, E. H. III (1997). Fires, hurricanes, and volcanoes: Comparing large disturbances. *BioScience*, 47, 758–768. <https://doi.org/10.2307/1313098>
- Turner, M. G., Gardner, R. H., & O'Neill, R. V. (2001). Landscape disturbance dynamics. In M. G. Turner, R. H. Gardner, & R. V. O'Neill (Eds.), *Landscape ecology in theory and practice* (pp. 157–199). New York, NY: Springer-Verlag.
- Turton, S. M. (1992). Undersorey light environments in a north-east Australian rain forest before and after a tropical cyclone. *Journal of Tropical Ecology*, 8, 241–252. <https://doi.org/10.1017/S0266467400006453>
- Turton, S. M. (2008). Landscape-scale impacts of Cyclone Larry on the forests of northeast Australia, including comparisons with previous cyclones impacting the region between 1858 and 2006. *Austral Ecology*, 33, 409–416. <https://doi.org/10.1111/j.1442-9993.2008.01896.x>
- Uriarte, M., Rivera, L. W., Zimmerman, J. K., Aide, T. M., Power, A. G., & Flecker, A. S. (2004). Effects of land use history on hurricane damage and recovery in a neotropical forest. *Plant Ecology*, 174, 49–58. <https://doi.org/10.1023/B:VEGE.0000046058.00019.d9>
- Van Bloem, S. J., Murphy, P. G., & Lugo, A. E. (2007). A link between hurricane-induced tree sprouting, high stem density and short canopy in tropical dry forest. *Tree Physiology*, 27, 475–480. <https://doi.org/10.1093/treephys/27.3.475>
- Venter, O., Sanderson, E. W., Magrath, A., Allan, J. R., Beher, J., Jones, K. R., ... Watson, J. E. M. (2016). Global terrestrial human footprint maps for 1993 and 2009. *Sci Data*, 3, 160067. <https://doi.org/10.1038/sdata.2016.67>
- Vincent, E. M., Lengaigne, M., Madec, G., Vialard, J., Samson, G., Jourdain, N. C., ... Jullien, S. (2012). Processes setting the characteristics of sea surface cooling induced by tropical cyclones. *Journal of Geophysical Research: Oceans*, 117, C02020.
- Vincent, E. M., Lengaigne, M., Vialard, J., Madec, G., Jourdain, N. C., & Masson, S. (2012). Assessing the oceanic control on the amplitude of sea surface cooling induced by tropical cyclones. *Journal of Geophysical Research: Oceans*, 117, C05023.
- Walsh, K. J. E., McBride, J. L., Klotzbach, P. J., Balachandran, S., Camargo, S. J., Holland, G., ... Sugi, M. (2016). Tropical cyclones and climate change. *Wiley Interdisciplinary Reviews: Climate Change*, 7, 65–89.
- Webb, E. L., Seamon, J. O., & Fa'umu, S. (2011). Frequent, low-amplitude disturbances drive high tree turnover rates on a remote, cyclone-prone Polynesian island. *Journal of Biogeography*, 38, 1240–1252. <https://doi.org/10.1111/j.1365-2699.2011.02505.x>
- Webb, E. L., Stanfield, B. J., & Jensen, M. L. (1999). Effects of topography on rainforest tree community structure and diversity in American Samoa, and implications for frugivore and nectarivore populations. *Journal of Biogeography*, 26, 887–897. <https://doi.org/10.1046/j.1365-2699.1999.00326.x>
- Whelan, R. J. (1995). *The ecology of fire*. Cambridge, UK: Cambridge University Press.
- Whigham, D., Dickinson, M., & Brokaw, N. (1999). Background canopy gap and catastrophic wind disturbances in tropical forests. In L. R. Walker (Ed.), *Ecosystems of the world 16: Ecosystems of disturbed ground* (pp. 223–252). Amsterdam, the Netherlands: Elsevier.
- White, P. S., & Jentsch, A. (2001). The search for generality in studies of disturbance and ecosystem dynamics. In K. Esser, U. Lüttge, J. W. Kadereit, & W. Beyschlag (Eds.), *Progress in botany: Genetics physiology systematics ecology* (pp. 399–450). Berlin, Heidelberg: Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-642-56849-7>
- White, P. S., & Pickett, S. T. A. (1985). Natural disturbance and patch dynamics: An introduction. In S. T. A. Pickett, & P. S. White (Eds.), *The ecology of natural disturbance and patch dynamics* (pp. 3–13). New York, NY: Academic Press.
- Whitmore, T. C. (1974). *Change with time and the role of cyclones in tropical rain forest on Kolombangara, Solomon Islands* by T.C. Whitmore. Oxford, UK: Commonwealth Forestry Institute, University of Oxford.
- Willoughby, H. E., Darling, R. W. R., & Rahn, M. E. (2006). Parametric representation of the primary hurricane vortex. Part II: A new family of sectionally continuous profiles. *Monthly Weather Review*, 134, 1102–1120. <https://doi.org/10.1175/MWR3106.1>
- Yamada, I. (1975). Forest ecological studies of the montane forest of Mt. Pangrango, West Java. *South East Asian Studies*, 13, 402–426.
- Yih, K., Boucher, D. H., Vandermeer, J. H., & Zamora, N. (1991). Recovery of the rain forest of southeastern Nicaragua after destruction by Hurricane Joan. *Biotropica*, 23, 106–113. <https://doi.org/10.2307/2388295>
- Yoo, J., Galewsky, J., Camargo, S. J., Korty, R., & Zamora, R. (2016). Dynamical downscaling of tropical cyclones from CCSM4 simulations of the Last Glacial Maximum. *Journal of Advances in Modeling Earth Systems*, 8, 1229–1247. <https://doi.org/10.1002/2016MS000685>
- Zeng, H., Chambers, J. Q., Negrón-Juárez, R. I., Hurr, G. C., Baker, D. B., & Powell, M. D. (2009). Impacts of tropical cyclones on U.S. forest tree mortality and carbon flux from 1851 to 2000. *Proceedings of*

- the National Academy of Sciences of the United States of America, 106, 7888–7892. <https://doi.org/10.1073/pnas.0808914106>
- Zhang, J., Nielsen, S. E., Mao, L., Chen, S., & Svenning, J.-C. (2016). Regional and historical factors supplement current climate in shaping global forest canopy height. *Journal of Ecology*, 104, 469–478. <https://doi.org/10.1111/1365-2745.12510>
- Zimmerman, J. K., Everham, E. M. III, Waide, R. B., Lodge, D. J., Taylor, C. M., & Brokaw, N. V. L. (1994). Responses of tree species to hurricane winds in subtropical wet forest in Puerto Rico: Implications for tropical tree life histories. *Journal of Ecology*, 82, 911–922. <https://doi.org/10.2307/2261454>
- ## DATA SOURCES
- Aiba, S., & Kitayama, K. (1999). Structure, composition and species diversity in an altitude-substrate matrix of rain forest tree communities on Mount Kinabalu, Borneo. *Plant Ecology*, 140, 139–157. <https://doi.org/10.1023/A:1009710618040>
- Blanc, L., Maury-Lechon, G., & Pascal, J. P. (2000). Structure, floristic composition and natural regeneration in the forests of Cat Tien National Park, Vietnam: An analysis of the successional trends. *Journal of Biogeography*, 27, 141–157. <https://doi.org/10.1046/j.1365-2699.2000.00347.x>
- Bongers, F., Popma, J., del Castillo, J. M., & Carabias, J. (1988). Structure and floristic composition of the lowland rain forest of Los Tuxtlas, Mexico. *Vegetatio*, 74, 55–80. <https://doi.org/10.1007/BF00045614>
- Bradford, M. G., Metcalfe, D. J., Ford, A., Liddell, M. J., & McKeown, A. (2014). Floristics, stand structure and aboveground biomass of a 25-ha rainforest plot in the wet tropics of Australia. *Journal of Tropical Forest Science*, 26, 543–553.
- Brearley, F. Q., Prajadinata, S., Kidd, P. S., Proctor, J., & Suriantata (2004). Structure and floristics of an old secondary rain forest in Central Kalimantan, Indonesia, and a comparison with adjacent primary forest. *Forest Ecology and Management*, 195, 385–397. <https://doi.org/10.1016/j.foreco.2004.02.048>
- Brewer, S. W., & Webb, M. A. H. (2002). A seasonal evergreen forest in Belize: Unusually high tree species richness for northern Central America. *Botanical Journal of the Linnean Society*, 138, 275–296. <https://doi.org/10.1046/j.1095-8339.2002.00016.x>
- Brockelman, W. Y., Nathalang, A., & Gale, G. A. (2011). The Mo Singto Forest Dynamics Plot, Khao Yai National Park, Thailand. *The Natural History Bulletin of the Siam Society*, 57, 35–55.
- Brokaw, N. V. L., Grear, J. S., Tripplett, K. J., Whitman, A. A., & Mallory, E. P. (1997). The Quebrada de Oro forest of Belize: Exceptional structure and high species richness. *Tropical Ecology*, 8, 247–258.
- Bunyavejchewin, S., Baker, P. J., LaFrankie, J. V., & Ashton, P. S. (2004). Huai Kha Khaeng forest dynamics plot, Thailand. In E. C. Losos, L. Leigh, & E. Giles (Eds.), *Tropical forest diversity and dynamism: Findings from a large-scale plot network* (pp. 585–598). Chicago, IL: Chicago University Press.
- Cadotte, M. W., Franck, R., Reza, L., & Lovett-Doust, J. (2002). Tree and shrub diversity and abundance in fragmented littoral forest of south-eastern Madagascar. *Biodiversity & Conservation*, 11, 1417–1436. <https://doi.org/10.1023/A:1016282023542>
- Clark, D. B., Hurtado, J., & Saatchi, S. S. (2015). Tropical rain forest structure, tree growth and dynamics along a 2700-m elevational transect in Costa Rica. *PLoS ONE*, 10, e0122905. <https://doi.org/10.1371/journal.pone.0122905>
- Condit, R., Aguilar, S., Hernandez, A., Perez, R., Lao, S., Angehr, G., ... Foster, R. B. (2004). Tropical forest dynamics across a rainfall gradient and the impact of an El Niño dry season. *Journal of Tropical Ecology*, 20, 51–72. <https://doi.org/10.1017/S0266467403001081>
- Culmsee, H., & Pitopang, R. (2009). Tree diversity in sub-montane and lower montane primary rain forests in Central Sulawesi. *Blumea*, 54, 119–123. <https://doi.org/10.3767/000651909X475473>
- Culmsee, H., Pitopang, R., Mangopo, H., & Sabir, S. (2011). Tree diversity and phytogeographical patterns of tropical high mountain rain forests in Central Sulawesi, Indonesia. *Biodiversity and Conservation*, 20, 1103–1123. <https://doi.org/10.1007/s10531-011-0019-y>
- D'Amico, C., & Gauthier, L. (2000). Inventory of a 1-ha lowland rainforest plot in Manongarivo (NW Madagascar). *Condollea*, 55, 319–340.
- De Gouvenain, R. C., & Silander, J. A. (2003). Do tropical storm regimes influence the structure of tropical lowland rain forests? *Biotropica*, 35, 166–180.
- DeWalt, S. J., Ickes, K., & James, A. (2016). Forest and community structure of tropical sub-montane rain forests on the island of Dominica, Lesser Antilles. *Caribbean Naturalist*, 1, 116–137.
- Fibich, P., Lepš, J., Novotný, V., Klimeš, P., Těšitel, J., Molem, K., ... Weiblen, G. D. (2016). Spatial patterns of tree species distribution in New Guinea primary and secondary lowland rain forest. *Journal of Vegetation Science*, 27, 328–339. <https://doi.org/10.1111/jvs.12363>
- Florens, F. B. V., Baider, C., Martin, G. M. N., & Strasberg, D. (2012). Surviving 370 years of human impact: What remains of tree diversity and structure of the lowland wet forests of oceanic island Mauritius? *Biodiversity and Conservation*, 21, 2139–2167. <https://doi.org/10.1007/s10531-012-0304-4>
- Franklin, J., Wiser, S. K., Drake, D. R., Burrows, L. E., & Sykes, W. R. (2006). Environment, disturbance history and rain forest composition across the islands of Tonga, Western Polynesia. *Journal of Vegetation Science*, 17, 233–244. <https://doi.org/10.1111/j.1654-1103.2006.tb02442.x>
- Gentry, A. H. (1995). Diversity and floristic composition of neotropical dry forests. In E. Medina, H. A. Mooney, & S. H. Bullock (Eds.), *Seasonally dry tropical forests* (pp. 146–194). Cambridge, UK: Cambridge University Press. <https://doi.org/10.1017/CBO9780511753398>
- Gillespie, T. W., Grijalva, A., & Farris, C. N. (2000). Diversity, composition, and structure of tropical dry forests in Central America. *Plant Ecology*, 147, 37–47. <https://doi.org/10.1023/A:1009848525399>
- Gillespie, T. W., Keppel, G., Pau, S., Price, J. P., Jaffre, T., & O'Neill, K. (2013). Scaling species richness and endemism of tropical dry forests on oceanic islands. *Diversity and Distributions*, 19, 896–906. <https://doi.org/10.1111/ddi.12036>
- Graham, A. W. (2006). The CSIRO rainforest permanent plots of North Queensland – Site, structural, floristic and edaphic descriptions, pp. 252. CSIRO and the Cooperative Research Centre for Tropical Rainforest Ecology and Management. Rainforest CRC, Cairns.
- Gunatilleke, C. V. S., Gunatilleke, I., Ashton, P. S., Ethugala, A. U. K., Weerasekera, N. S., & Esufali, S. (2004). Sinharaja forest dynamics plot, Sri Lanka. *Tropical forest diversity and dynamism: Findings from a large-scale plot network* (pp. 599–608).
- Hamann, A., Barbon, E. B., Curio, E., & Madulid, D. A. (1999). A botanical inventory of a submontane tropical rainforest on Negros Island, Philippines. *Biodiversity and Conservation*, 8, 1017–1031. <https://doi.org/10.1023/A:1008847704539>
- Ibanez, T., Blanchard, E., Hequet, V., Keppel, G., Laidlaw, M., Pouteau, R., ... Birnbaum, P. (2017). High endemism and stem density distinguish New Caledonian from other high-diversity rainforests in the Southwest Pacific. *Annals of Botany*, 121, 25–35.
- Ibañez, A., & Castroviejo, S. (2005). The vegetation of Bahia Honda Region (Veraguas, Panama). In S. Castroviejo & A. Ibañez (Eds.), *Studies of the biodiversity of the Bahia Honduras Region (Veraguas, Panama)*. Madrid, Spain: Consejo superior de investigaciones científicas.
- Jaffré, T., & Veillon, J. M. (1991). Etude floristique et structurale de deux forêts denses humides sur roches ultrabasiqes en Nouvelle-Calédonie. *Bulletin du Muséum National d'Histoire Naturelle Section B, Adansonia*, 4^e sér., 12, 243–273.
- Jaffré, T., & Veillon, J. M. (1995). Structural and floristic characteristics of a rain forest on schist in New Caledonia: A comparaison with an ultramafic rain forest. *Bulletin du Muséum National d'Histoire Naturelle Section B, Adansonia*, 4^e sér., 17, 201–226.

- Johnston, M., & Gillman, M. (1995). Tree population studies in low-diversity forests, Guyana. I. Floristic composition and stand structure. *Biodiversity & Conservation*, 4, 339–362. <https://doi.org/10.1007/BF00058421>
- Josse, C., & Balslev, H. (1994). The composition and structure of a dry, semideciduous forest in western Ecuador. *Nordic Journal of Botany*, 14, 425–434. <https://doi.org/10.1111/j.1756-1051.1994.tb00628.x>
- Kanzaki, M., Hara, M., Yamakura, T., Ohkubo, T., Tamura, N., Sringeriyuang, K., ... Bunyavejchewin, S. (2004). Doi Inthanon forest dynamics plot, Thailand. In E. C. Losos, L. Leigh & E. Giles (Eds.), *Tropical forest diversity and dynamism: Findings from a large-scale plot network* (pp. 585–598). Chicago, IL: Chicago University Press.
- Kartawinata, K., Partomihardjo, P. T., Yusuf, R., Abdulhadi, R., & Riswan, S. (2008). Floristics and structure of a lowland dipterocarp forest at Wanariset Samboja, East Kalimantan, Indonesia. *Reinwardtia*, 12, 301–323.
- Kartawinata, K., Samsudin, I., Heriyanto, M., & Afriastini, J. J. (2004). A tree species inventory in a one-hectare plot at the Batang Gadis National Park, North Sumatra, Indonesia. *Reinwardtia*, 12, 145–147.
- Kearsley, E., Moonen, P. C., Hufkens, K., Doetterl, S., Lisingo, J., Boyemba Bosela, F., ... Verbeeck, H. (2017). Model performance of tree height-diameter relationships in the central Congo Basin. *Annals of Forest Science*, 74, 7. <https://doi.org/10.1007/s13595-016-0611-0>
- Kelly, D. L., Tanner, E. V. J., Kapos, V., Dickinson, T. A., Goodfriend, G. A., & Fairbairn, P. A. (1988). Jamaican limestone forests: Floristics, structure and environment of three examples along a rainfall gradient. *Journal of Tropical Ecology*, 4, 121–156. <https://doi.org/10.1017/S0266467400002649>
- Keppel, G., Buckley, Y. M., & Possingham, H. P. (2010). Drivers of lowland rain forest community assembly, species diversity and forest structure on islands in the tropical South Pacific. *Journal of Ecology*, 98, 87–95. <https://doi.org/10.1111/j.1365-2745.2009.01595.x>
- Kitamura, S., Suzuki, S., Yumoto, T., Chuailua, P., Plongmai, K., Poonswad, P., ... Suckasam, C. (2005). A botanical inventory of a tropical seasonal forest in Khao Yai National Park, Thailand: Implications for fruit-frugivore interactions. *Biodiversity and Conservation*, 14, 1241–1262. <https://doi.org/10.1007/s10531-004-7848-x>
- Kohira, M., Ninomiya, I., Ibrahim, A. Z., & Latiff, A. (2001). Diversity, diameter structure and spatial pattern of trees in a semi-evergreen rain forest on Langkawi Island, Malaysia. *Journal of Tropical Forest Science*, 13, 460–476.
- Laidlaw, M., Kitching, R., Goodall, K., Small, A., & Stork, N. (2007). Temporal and spatial variation in an Australian tropical rainforest. *Austral Ecology*, 32, 10–20. <https://doi.org/10.1111/j.1442-9993.2007.01739.x>
- Lee, H. S., Tan, S., Davies, S. J., LaFrankie, J. V., Ashton, P. S., Yamakura, T., ... Harrison, R. (2004). Lambir forest dynamics plot, Sarawak, Malaysia. In E. C. Losos, L. Leigh, & E. Giles (Eds.), *Tropical forest diversity and dynamism: Findings from a large-scale plot network* (pp. 585–598). Chicago, IL: Chicago University Press.
- Lu, X. T., Yin, J. X., & Tang, J. W. (2010). Structure, tree species diversity and composition of tropical seasonal rainforests in Xishuangbanna, South-West China. *Journal of Tropical Forest Science*, 22, 260–270.
- Lu, X., Zang, R., Ding, Y., Letcher, S. G., Long, W., & Huang, Y. (2014). Variations and trade-offs in functional traits of tree seedlings during secondary succession in a tropical lowland rain forest. *Biotropica*, 46, 404–414. <https://doi.org/10.1111/btp.12125>
- Makana, J.-R., Hart, T. B., Liengola, I., Ewango, C., Hart, J. A., & Condit, R. Ituri forest dynamics plots, Democratic Republic of Congo.
- Manokaran, N., Quah, E. S., Ashton, P. S., LaFrankie, J. V., Noor, N. S. M., Ahmad, W. M. S. W., & Okuda, T. (2004). Pasoh forest dynamics plot, Peninsular Malaysia. In E. C. Losos, L. Leigh, & E. Giles (Eds.), *Tropical forest diversity and dynamism: Findings from a large-scale plot network* (pp. 585–598). Chicago, IL: Chicago University Press.
- Martin, P. H., Sherman, R. E., & Fahey, T. J. (2004). Forty years of tropical forest recovery from agriculture: Structure and floristics of secondary and old-growth riparian forests in the Dominican Republic. *Biotropica*, 36, 297–317.
- Nesheim, I., Halvorsen, R., & Nordan, I. (2010). Plant composition in the Maya Biosphere Reserve: Natural and anthropogenic influences. *Plant Ecology*, 208, 93–122. <https://doi.org/10.1007/s11258-009-9691-3>
- Newbery, D. M., Gartlan, J. S., McKey, D. B., & Waterman, P. G. (1986). The influence of drainage and soil phosphorus on the vegetation of Douala-Edea Forest Reserve, Cameroun. *Vegetatio*, 65, 149–162. <https://doi.org/10.1007/BF00044815>
- Okali, D. U. U., & Ola-Adams, B. A. (1987). Tree population changes in treated rain forest at Omo Forest Reserve, south-western Nigeria. *Journal of Tropical Ecology*, 3, 291–313. <https://doi.org/10.1017/S0266467400002297>
- Parthasarathy, N., & Karthikeyan, R. (1997). Plant biodiversity inventory and conservation of two tropical dry evergreen forests on the Coromandel coast, south India. *Biodiversity and Conservation*, 6, 1063–1083. <https://doi.org/10.1023/A:1018328016810>
- Pascal, J.-P., & Pelissier, R. (1996). Structure and floristic composition of a tropical evergreen forest in South-West India. *Journal of Tropical Ecology*, 12, 191–214. <https://doi.org/10.1017/S026646740000941X>
- Pereira, I. M., Andrade, L. A., Sampaio, E. S. B. V., & Barbosa, M. R. V. (2003). Use-history effects on structure and flora of Caatinga. *Biotropica*, 35, 154–165.
- Phillips, O. L., Hall, P., Gentry, A. H., Sawyer, S. A., & Vásquez, R. (1994). Dynamics and species richness of tropical rain forests. *Proceedings of the National Academy of Sciences of the United States of America*, 91, 2805–2809. <https://doi.org/10.1073/pnas.91.7.2805>
- Phillips, O., & Miller, J. S. (2002). *Global patterns of plant diversity: Alwyn H. Gentry's forest transect data set*. St. Louis, MO: MBG Press, Missouri Botanical Garden.
- Poulsen, A. D., Nielsen, I. C., Tan, S., & Balslev, H. (1996). A quantitative inventory of trees in one hectare mixed dipterocarp forest in Temburong, Brunei Darussalam. *Tropical Rainforest Research*, 74, 139–150. <https://doi.org/10.1007/978-94-009-1685-2>
- Proctor, J., Anderson, J. M., Chai, P., & Vallack, H. W. (1983). Ecological studies in 4 contrasting lowland rain forests in Gunung-Mulu-National-Park, Sarawak. 1. Forest environment, structure and floristics. *Journal of Ecology*, 71, 237–260. <https://doi.org/10.2307/2259975>
- Proctor, J., Baker, A. J. M., Van Balgooy, M. M. J., Bruijnzeel, L. A., Jones, S. H., & Madulid, D. A. (2000). Mount Bloomfield, Palawan, Philippines: Forests on greywacke and serpentinized peridotite. *Edinburgh Journal of Botany*, 57, 121–139. <https://doi.org/10.1017/S0960428600000081>
- Proctor, J., Lee, Y. F., Langley, A. M., Munro, W. R. C., & Nelson, T. (1988). Ecological studies on Gunung Silam, a small ultrabasic mountain in Sabah, Malaysia. I. Environment, forest structure and floristics. *Journal of Ecology*, 76, 320–340. <https://doi.org/10.2307/2260596>
- Quigley, M. F., & Platt, W. J. (2003). Composition and structure of seasonally deciduous forests in the Americas. *Ecological Monographs*, 73, 87–106. [https://doi.org/10.1890/0012-9615\(2003\)073\[0087:CASOSD\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2003)073[0087:CASOSD]2.0.CO;2)
- Rai, S. N., & Proctor, J. (1986). Ecological studies on four rainforests in Karnataka, India: I. Environment, structure, floristics and biomass. *Journal of Ecology*, 74, 439–454. <https://doi.org/10.2307/2260266>
- Rajkumar, M., & Parthasarathy, N. (2008). Tree diversity and structure of Andaman giant evergreen forests, India. *Taiwania*, 53, 356–358.
- Rakotomalaza, P. J., & Messmer, N. (1999). Structure and floristic composition of the vegetation in the Réserve Naturelle Intégrale d'Andohahela, Madagascar. In S. M. Goodman (Ed.), *A floral and faunal inventory of the Réserve Naturelle Intégrale d'Andohahela, Madagascar: With reference to elevational variation* (pp. 51–96). Chicago, IL: Field Museum of Natural History.
- Rasingam, L., & Parthasarathy, N. (2009). Tree species diversity and population structure across major forest formations and disturbance categories in Little Andaman Island, India. *Tropical Ecology*, 50, 89–102.

- Sheil, D., Kartawinata, K., Samsudin, I., Priyadi, H., & Afriastini, J. J. (2010). The lowland forest tree community in Malinau, Kalimantan (Indonesian Borneo): results from a one-hectare plot.
- Sist, P., & Saridan, A. (1999). Stand structure and floristic composition of a primary lowland dipterocarp forest in East Kalimantan. *Journal of Tropical Forest Science*, 11, 1–11.
- Small, A., Martin, T. G., Kitching, R. L., & Wong, K. M. (2004). Contribution of tree species to the biodiversity of a 1 ha Old World rainforest in Brunei, Borneo. *Biodiversity and Conservation*, 13, 2067–2088. <https://doi.org/10.1023/B:BIOC.0000040001.72686.e8>
- Strasberg, D. (1996). Diversity, size composition and spatial aggregation among trees on a 1-ha rain forest plot at La Reunion. *Biodiversity and Conservation*, 5, 825–840. <https://doi.org/10.1007/BF00054737>
- Su, S.-H., Chang-Yang, C.-L., Lu, C.-L., Tsui, C.-C., Lin, T.-T., Chiou, W.-L., ... Hsieh, C.-F. (2007). Fushan subtropical forest dynamics plot: Tree species characteristics and distribution patterns. *TFRI extension* (p. 285). Taiwan Forestry Research Institute.
- Sudhakar Reddy, C., Babar, S., Amarnath, G., & Pattanaik, C. (2011). Structure and floristic composition of tree stand in tropical forest in the Eastern Ghats of northern Andhra Pradesh, India. *Journal of Forestry Research*, 22, 491–500. <https://doi.org/10.1007/s11676-011-0193-5>
- Sukardjo, S., Hagihara, A., Yamakura, T., & Ogawa, H. (1990). Floristic composition of a tropical rain forest in Indonesian Borneo. *Bulletin of the Nagoya University Forests*, 10, 1–44.
- Sukumar, R., Suresh, H. S., Dattaraja, H. S., John, R., & Joshi, N. V. (2004). Mudumalai forest dynamics plot, India. In E. C. Losos, L. Leigh, & E. Giles (Eds.), *Tropical forest diversity and dynamism: Findings from a large-scale plot network* (pp. 585–598). Chicago, IL: Chicago University Press.
- Sun, I.-F., & Hsieh, C.-F. (2004). Nanjenshan forest dynamics plot, Taiwan. In E. C. Losos, L. Leigh, & E. Giles (Eds.), *Tropical forest diversity and dynamism: Findings from a large-scale plot network* (pp. 585–598). Chicago, IL: Chicago University Press.
- Sussman, R. W., & Rakotozafy, A. (1994). Plant diversity and structural analysis of a tropical dry forest in Southwestern Madagascar. *Biotropica*, 26, 241–254. <https://doi.org/10.2307/2388845>
- Thomas, D. W., Kenfack, D., Chuyong, G. B., Moses, S. N., Losos, E. C., Condit, R. S., & Songwe, N. C. (2003). Tree species of south-western Cameroon: Tree distribution maps, diameter tables, and species documentation of the 50-hectare Korup Forest Dynamics Plot. Washington, DC: Center for Tropical Forest Science of the Smithsonian Tropical Research Institute and Bioresources Development and Conservation Programme-Cameroon.
- Thompson, J., Brokaw, N., Zimmerman, J. K., Waide, R. B., Everham, E. M., Lodge, D. J., ... Fluett, M. (2002). Land use history, environment, and tree composition in a tropical forest. *Ecological Applications*, 12, 1344–1363. [https://doi.org/10.1890/1051-0761\(2002\)012\[1344:LUHEAT\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2002)012[1344:LUHEAT]2.0.CO;2)
- Thompson, J., Brokaw, N., Zimmerman, J. K., Waide, R. B., Everham III, E. M., & Schaefer, D. A. Luquillo forest dynamics plot, Puerto Rico, United States.
- Thompson, J., Proctor, J., Viana, V., Milliken, W., Ratter, J. A., & Scott, D. A. (1992). Ecological studies on a lowland evergreen rain forest on Maraca Island, Roraima, Brazil. I. Physical environment, forest structure and leaf chemistry. *Journal of Ecology*, 80, 689–703. <https://doi.org/10.2307/2260860>
- Valentia, R., Condit, R., Foster, R. B., Romoleroux, K., Villa Munoz, G., Svenning, J.-C., ... Balslev, H. Yasuni forest dynamic plot, Ecuador.
- Webb, E. L., & Fa'aumu, S. (1999). Diversity and structure of tropical rain forest of Tutuila, American Samoa: Effects of site age and substrate. *Plant Ecology*, 144, 257–274. <https://doi.org/10.1023/A:1009862618250>
- Weiblen, G. D. (1998). Composition and structure of a one hectare forest plot in the Crater Mountain Wildlife Management Area, Papua New Guinea. *Science in New Guinea*, 24, 23–32.
- Whitfield, T. J. S., Lasky, J. R., Damas, K., Sosanika, G., Molem, K., & Montgomery, R. A. (2014). Species richness, forest structure, and functional diversity during succession in the New Guinea lowlands. *Biotropica*, 46, 538–548. <https://doi.org/10.1111/btp.12136>
- Whitmore, T. C., & Sidiyasa, K. (1986). Composition and structure of a lowland rain forest at Toraut, Northern Sulawesi. *Kew Bulletin*, 41, 747–757. <https://doi.org/10.2307/4103127>
- Whitmore, T. C., Sidiyasa, K., & Whitmore, T. J. (1987). Tree species enumeration of 0.5 hectare on Halmahera. *The Gardens' Bulletin, Singapore*, 40, 31–34.
- Wilkie, P., Argent, G., Cambell, E., & Saridan, A. (2004). The diversity of 15 ha of lowland mixed dipterocarp forest, Central Kalimantan. *Biodiversity & Conservation*, 13, 695–708. <https://doi.org/10.1023/B:BIOC.0000011721.04879.79>
- Wright, D. D., Jessen, J. H., Burke, P., & de Silva Garza, H. G. (1997). Tree and liana enumeration and diversity on a one-hectare plot in Papua New Guinea. *Biotropica*, 29, 250–260. <https://doi.org/10.1111/j.1744-7429.1997.tb00426.x>
- Zent, E. L., & Zent, S. (2004). Floristic composition, structure, and diversity of four forest plots in the Sierra Maigualida, Venezuelan Guayana. *Biodiversity and Conservation*, 13, 2453–2484. <https://doi.org/10.1023/B:BIOC.0000048447.40238.f2>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Ibanez T, Keppel G, Menkes C, et al.

Globally consistent impact of tropical cyclones on the structure of tropical and subtropical forests. *J Ecol*.

2019;107:279–292. <https://doi.org/10.1111/1365-2745.13039>